

§ 201.316

ent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult physician promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by physician.”

§201.316 Drugs with thyroid hormone activity for human use; required warning.

(a) Drugs with thyroid hormone activity have been promoted for, and continue to be dispensed and prescribed for, use in the treatment of obesity, although their safety and effectiveness for that use have never been established.

(b) Drugs for human use with thyroid hormone activity are misbranded within the meaning of section 502 of the Federal Food, Drug, and Cosmetic Act unless their labeling bears the following boxed warning at the beginning of the “Warnings” section:

Drugs with thyroid hormone activity, alone or together with other therapeutic agents, have been used for the treatment of obesity. In euthyroid patients, doses within the range of daily hormonal requirements are ineffective for weight reduction. Larger doses may produce serious or even life-threatening manifestations of toxicity, particularly when given in association with sympathomimetic amines such as those used for their anorectic effects.

[43 FR 22009, May 23, 1978]

§201.317 Digitalis and related cardiotonic drugs for human use in oral dosage forms; required warning.

(a) Digitalis and related cardiotonic drugs for human use in oral dosage forms have been promoted for, and continue to be dispensed and prescribed for, use in the treatment of obesity, although their safety and effectiveness for that use have never been established.

(b) Digitalis and related cardiotonic drugs for human use in oral dosage forms are misbranded within the mean-

21 CFR Ch. I (4–1–96 Edition)

ing of section 502 of the Federal Food, Drug, and Cosmetic Act unless their labeling bears the following boxed warning at the beginning of the “Warnings” section:

Digitalis alone or with other drugs has been used in the treatment of obesity. This use of digoxin or other digitalis glycosides is unwarranted. Moreover, since they may cause potentially fatal arrhythmias or other adverse effects, the use of these drugs in the treatment of obesity is dangerous.

(c) This section does not apply to digoxin products for oral use, which shall be labeled according to the requirements of §310.500 of this chapter.

[43 FR 22009, May 23, 1978]

§201.319 Water-soluble gums, hydrophilic gums, and hydrophilic mucilloids (including, but not limited to agar, alginic acid, calcium polycarbophil, carboxymethylcellulose sodium, carrageenan, chondrus, glucomannan ((B-1,4 linked) polymannose acetate), guar gum, karaya gum, kelp, methylcellulose, plantago seed (psyllium), polycarbophil, tragacanth, and xanthan gum) as active ingredients; required warnings and directions.

(a) Reports in the medical literature and data accumulated by the Food and Drug Administration indicate that esophageal obstruction and asphyxiation have been associated with the ingestion of water-soluble gums, hydrophilic gums, and hydrophilic mucilloids including, but not limited to, agar, alginic acid, calcium polycarbophil, carboxymethylcellulose sodium, carrageenan, chondrus, glucomannan ((B-1,4 linked) polymannose acetate), guar gum, karaya gum, kelp, methylcellulose, plantago seed (psyllium), polycarbophil, tragacanth, and xanthan gum. Esophageal obstruction and asphyxiation due to orally-administered drug products containing water-soluble gums, hydrophilic gums, and hydrophylic mucilloids as active ingredients are significant health risks when these products are taken without